

# Study on Invasive Procedures of the Chest Lesions: Are they must be Performed and Why?

Hilendarov A<sup>1\*</sup>, Georgiev A<sup>1</sup>, Chervenkov A<sup>1</sup>

<sup>1</sup>Medical University-Plovdiv, Medical Faculty, Department of Diagnostic Imaging, Plovdiv, Bulgaria.

**\*Correspondence:** Dr. Atanas D. Hilendarov, Associate Professor, Dept of Diagnostic Imaging, MU Plovdiv 49 Volga str, ap.14, 6 fl, 4002 Plovdiv, Bulgaria.

**Received:** September 09, 2026 | **Published:** October 01, 2026

## How to cite this article:

Hilendarov A. Study on Invasive Procedures of the Chest Lesions: Are they must be Performed and Why? *J Case Rep Rev Med.* 2026;2(4):1-5. doi:10.64978/jcrrm.2026.10010133.

## Abstract

**Introduction:** Chest tumours, in particular lung cancer, remain one of the most common causes of death worldwide. Using MDCT, an increasing number of lung and mediastinal lesions is detected and histological diagnosis is often necessary to determine the most appropriate management of these lesions.

**Aims:** At this article we have the aims to describe the invasive procedures of the chest lesions - indications, contraindications, technical aspects, and the diagnostic accuracy of the percutaneous lung biopsies.

**Methods:** Fine-needle aspiration biopsy (FNAB) and core-needle biopsy (CNB) currently is the predominant method for obtaining tissue specimens in patients with lung lesions. In many cases, treatment protocols are based on histological information. In 85 of all 97 patients included in ours study FNAB biopsy is performed and in 12 CNB, when technically feasible, or in cases where other techniques (such as bronchoscopy with lavage) are inconclusive. The 19-22G disposable needles were used.

**Results:** In all 76 patients, aged 21-79 years with pulmonary lesions with dimensions of 2.0 cm or less FNB under CT control are performed. In 13 cases FNA under US control is performed due to the superficial localization of the lesions. Cytological and \ evaluation of FNAB samples were performed in all patients. Diagnostic sensitivity and accuracy are calculated. Assess the type of complications that occurred. FNAB and CNB, with the latter, demonstrated to have a slightly higher overall sensitivity, specificity and accuracy.

**Conclusion:** Percutaneous FNAB and CNB are safe procedures even though a few complications are possible: pneumothorax, pulmonary haemorrhage is common, while air embolism and seeding are rare, but with severe consequences.

**Keywords:** core-needle biopsy, diagnostic accuracy, fine-needle aspiration biopsy, lung lesion

## Introduction

Lung cancers constitute a large group of diseases as far as their histopathology is concerned. The method of treatment depends on the lesion type. Nowadays, there are two basic groups of cancers identified on the basis of the treatment method: small cell cancers and non-small cell cancers. The first type is characterised by a rapid dissemination, so that in most of the cases, the cancer is highly advanced at the moment of the diagnosis. The treatment of choice is chemotherapy. Non-small cell cancers constitute a group of tumours with heterogeneous histopathology. However, their treatment methods are similar and depend to a larger

degree on the clinical advancement of the cancer rather than on its histopathological structure. The treatment of choice in case of relatively unadvanced non-small cell cancers is the surgical procedure extended by chemotherapy or radiotherapy. In case of more advanced non-small cell cancers, the therapeutic process may join the induction chemotherapy with a surgical procedure or apply chemo- or radiotherapy only . Taking into consideration all these factors, it is extremely important to obtain the histopathological diagnosis.

CT-guided fine-needle biopsy (FNB) of focal lung lesions is one of the possibilities of obtaining histopathological diagnosis in

pulmonary diseases. Its place in the algorithm is determined by the invasiveness. In case of no diagnosis after bronchoscopy or endobronchial ultrasonography (EBUS) guided biopsy, CT-guided FNB can become an alternative for more invasive procedures, such as open lung biopsy – thoracotomy.

Chest tumors, in particular lung cancer, remain one of the most common causes of death worldwide. Using MDCT, an increasing number of lung and mediastinal lesions is detected and histological diagnosis is often necessary to determine the most appropriate management of these lesions.<sup>1</sup> Various imaging techniques including computed tomography (CT) fluoroscopy and ultrasound (US) can be used to guide chest biopsies. CT is the most frequently used methods because of its high spatial and contrast resolution.

Indications of imaging-guided chest biopsy have significantly changed as a result of technical advances in needle types, imaging modalities, pathological analysis and immunohistological techniques. At present, the growing list of indications includes histological diagnosis of undetermined and otherwise not characteristic pulmonary, mediastinal and chest wall lesions as well as biopsy or re-biopsy of known malignant lesions to obtain histological material for targeted therapy.<sup>2</sup>

The use of MDCT has increased with the detection of small pulmonary nodules. In many patients, pulmonary nodules are incidentally detected in the general population<sup>3</sup> in patients without clinical manifestation.

Park and others present their results of total of 15,809 chest radiographs were collected from two tertiary hospitals (7204 normal and 8605 abnormal with nodule/mass, interstitial opacity, pleural effusion, or pneumothorax). Except for the test set (100 normal and 100 abnormal (nodule/mass, 70; interstitial opacity, 10; pleural effusion, 10; pneumothorax, 10)), radiographs were used to develop a DLD system for detecting multiclass lesions. The diagnostic performance of the developed model and that of nine observers with varying experiences were evaluated and compared using area under the receiver operating characteristic curve (AUROC), on a per-image basis, and jackknife alternative free-response receiver operating characteristic figure of merit (FOM) on a per-lesion basis. The false-positive fraction was also calculated. The DLD system showed the potential for detection of lesions and pattern classification on chest radiographs,<sup>3</sup> performing normal/abnormal classifications and achieving high diagnostic performance.

If a nodule is initially identified at conventional chest radiography, CT investigation is necessary to establish the lesion, estimate the malignancy<sup>3-6</sup> and identify lymphadenopathies or other accessible sites for biopsy - extra-thoracic metastases.<sup>7</sup> PET-CT examination may also be helpful in the investigation of pulmonary nodules, reducing the need for puncture of non-enhancing solid nodules.<sup>8</sup> It should be borne in mind that active inflammation (i.e. active tuberculosis, histoplasmosis, rheumatoid nodules) may cause false positives on PET/CT scans because of their high glucose metabolism.<sup>9</sup> Although low-grade malignant tumours, such as carcinoid<sup>10</sup> or low-grade adenocarcinoma,<sup>11</sup> may produce false-negative results because of their low glucose metabolism. In these patients, follow-up with CT must be performed to demonstrate regression of the nodule after therapy or if the increase in size is evident.<sup>12</sup>

Mediastinal masses are most frequently located in the anterior mediastinum and include a variety of different entities, such as thymic malignancy, lymphomas, endocrine tumours and malignant

germ cell tumours. In the absence of typical clinical and imaging features, histological diagnosis is necessary.

## 2. Material and Methods

Fine-needle aspiration biopsy (FNAB) and core-needle biopsy (CNB) currently is the predominant method for obtaining tissue specimens in patients with lung lesions. In many cases treatment protocols are based on histological information. In 85 of all 97 patients included in our study, performed at the University Hospital in Dept of Diagnostic Imaging, FNAB biopsy is performed and in 12 patients (CNB), when technically feasible, or in cases where other techniques (such as bronchoscopy with lavage) are inconclusive. The 19-22G disposable needles were used.

Imaging guidance methods we used to guide chest biopsies. We preferred CT guidance (Fig. 1).<sup>13-17</sup> Parameters affecting the selection of the most appropriate imaging technique are the size and visibility of the lesion as well as its relationship with critical anatomical structures. Whenever possible, chest biopsies should be performed under US guidance (Fig. 2) to use the advantages of real-time monitoring without radiation to patients and operators.<sup>18</sup>

US guidance is limited to superficial lesions adjacent to the chest wall or to lesions delineated by pleural effusion sufficient to create a suitable interface for ultrasound penetration. In the significant majority of cases, MDCT is the preferred guidance method for chest biopsies due to its optimal spatial and contrast resolution. An intravenous contrast agent can be used to differentiate target lesions from atelectasis, necrosis and vascular structures.

The patient should be positioned prone, supine or lying on the side, based on the previously planned access site. Whenever possible, the needle access site should be cephalic to the ribs to avoid intercostal vessel and nerves puncture. The skin in the access site should be sterilised with an antiseptic solution and subcutaneous tissues should be infiltrated with lidocaine. Breath-hold capabilities are extremely different. It is easier to target larger tumours (>2/3 cm) instructing the patient to breath freely with shallow respiration. In the remaining cases, the patient can be instructed to maintain an inspiratory or expiratory apnoea to allow easier access to target lesions.

Needle choice is based on the size and location of the lesion, planned needle pathway, status of the lung parenchyma and operator preference. Some of the used fine-needle aspiration devices include Chiba, Westcott, Greene (Cook) and Turner (Cook) needles, which have circumferentially sharpened tips allowing for sampling.

Core biopsy needles are designed to collect a small piece of tissue intended for surgical pathology analysis and can be designed as end-cutting or side-cutting devices.

The most commonly used core biopsy needle is the Tru-Cut, which consists of an outer cutting cannula and an inner slotted stylet. The presence of an on-site pathologist may reduce the number of biopsies needed.

## 3. Results

In 76 patients, aged 21-79 years with pulmonary lesions with dimensions of 2.0 cm or less FNB under CT control are performed. In 13 cases FNA under US control are performed due to the superficial localization of the lesions. Cytological and histological evaluation of FNAB samples were performed in all patients.

Diagnostic sensitivity and accuracy are calculated. Assess the number and type of complications that occurred. Fine-needle aspiration biopsy (FNAB) and core-needle biopsy (CNB), with the latter demonstrated to have a slightly higher overall sensitivity, specificity and accuracy.

We prefer MDCT for the guidance of FNB due to the better biopsy needle path control and the visibility of small-sized and deeply located lung lesions. We have a wide variation in diagnostic accuracies of fine-needle aspiration biopsy (FNAB) ranging from 64% to 97%.




Fig. 1: CT guided biopsy. Indications for FNAB (a) Mass lesion following negative bronchoscopy (b) CNB of unknown mass in the anterior part of mediastinum.



Fig. 2: Ultrasound-guided FNB biopsy of a pulmonary nodule in the left lower lobe. The needle was 20 G due to suspicions of liquide nature of the lesion




Fig. 3: CT image presenting left pulmonary lobe . CT-guided Core chest biopsy of a pulmonary nodule in the left lower lobe.




Fig. 4: CT-guided chest biopsy (a) Axial CT image before the procedure showing the left subpleural nodule; (b) Post-procedural axial CT image showing small PNX (arrows) as a complication of transthoracic




Fig. 5: CT-guided chest biopsy of a pulmonary nodule (a) Axial CT image showing a pulmonary nodule in the right lower lobe; (b) Postprocedural CT image demonstrates perilesional haemorrhage as ground-glass opacity around the nodule (arrows).

In our patients, the effects of lesion location on the diagnostic accuracy of imaging-guided biopsy in chest tumours were evaluated.

#### 4. Discussion

In literature reports the sensitivity in diagnosing lung tumours decreased from 100% in peripheral lesions to 82% in central lesions .Wang et al.<sup>17,18</sup> recently compared the rates of complications and diagnostic accuracy of CT-guided biopsy in peripheral versus paramediastinal lesions, demonstrating how this technique may be safe and reliable even for deeply located tumours; in particular, their paper reports diagnostic accuracy of 95.4% in paramediastinal lesions and 94.7% in peripheral lesions with a sensitivity of 95.6% and 94.2% respectively.

Once the biopsy has been performed, we perform control by CT of the chest is obtained to identify any immediate post-procedural complications. According to some authors, the patients should be rolled over onto the punctured side to reduce the risk of delayed PNX; Chest films are usually acquired after 4 h to detect possible asymptomatic PNX. If the clinical suspicion of a PNX arises, chest radiography must be obtained immediately.

Pneumothorax (PNX) is the most common complication after imaging-guided chest biopsy; it is most frequently detected after lung biopsy but can also occur even after biopsies of mediastinal, pleural and chest wall lesions. PNX occurs during or immediately after the procedure and can be detected on postprocedural control scans (Fig.4). The incidence of PNX has been reported to be up to 61% with an average risk of 20% . Risk factors for PNX can be related to patient or lesions features, but also the biopsy technique. In particular, the rate of PNX increases with the patient age and severity of underlying lung disease (e.g. emphysema or chronic obstructive disease) as well as in smaller and deeper lesions. Technical risk factors include the type and size of the biopsy needle, longer procedure duration, biopsies in the middle or lower lobe, transgression of a fissure and multiple needle repositioning

or pleural passes. A PNx developed during the procedure can be immediately aspirated through the introducer needle or a separate needle inserted into the pleural space, preventing further enlargement; however, some authors suggest placing a chest tube if aspirated air is greater than 670 ml. PNx can be significant (>30% of lung volume), increase over time or become symptomatic. In these cases, small-calibre, 6- to 9-French catheters can be safely and easily placed.

Haemorrhage represents the second most common complication after imaging-guided biopsy. PH may occur with or without haemoptysis and can be easily detected on screening post-biopsy CT scan as a perilesional or needle tract ground-glass opacity. (Fig.5). The occurrence rates of PH are estimated to be from 4 to 27% (with an average incidence of 11%), while haemoptysis risk is up to 5. Complication does not need any treatment and the only recommendations are to place the patient in a lateral position, with the biopsy side down, to avoid aspiration of blood into the unaffected lung. Occasionally a larger, higher-grade PH occurs and pro-coagulative treatment may be needed. Risk factors for higher-grade PH include older patients with emphysema, pulmonary hypertension, coaxial technique, subsolid lesions, and lesion size smaller than 3 cm.<sup>19,20</sup>

The occurrence of systemic air embolism (SAE) in the left atrium, left ventricle or systemic circulation is rare (incidences between 0.01% and 0.21%), but potentially fatal cardiac infarct.<sup>21</sup> There are three mechanisms in particular responsible for SAE during biopsy: placement of the needle tip in a pulmonary vein, formation of a bronchial-venous or alveolar-venous fistula and opening the outer cannula of a coaxial biopsy needle to the atmosphere.

Tumour seeding through the needle track represents a very rare complication with a prevalence reported in the literature between 0.012 and 0.061%. The real clinical relevance is still discussed, but it is obvious that tumour seeding along the needle track can significantly change patient management and life expectancy and should be strictly avoided. Tumour seeding is reported to be more frequently observed after imaging-guided core needle biopsy of pleural mesothelioma.

## 5. Conclusion

In conclusion, chest biopsy under imaging control is an interventional procedure very important for some clinical conditions of pneumatological, oncological and surgical interest. Fine-needle biopsy of the focal lung lesions is an effective and a relatively safe method, which can replace the more invasive diagnostic thoracotomy in the majority of patients.

This procedure may appear very simple, but radiologists approaching it for the first time must consider several clinical and technical variables significantly affecting the final results, in terms of both diagnostic accuracy and patients' safety.

There are no ethical issues associated with this manuscript.

## References

- Viggiano RW, Swensen SJ, Rosenow EC 3rd. Evaluation and management of solitary and multiple pulmonary nodules. *Clin Chest Med*. 1992;13(1):83-95. doi:10.1016/S0272-5231(21)00598-4.
- Hammerschlag G, Cao J, Gumm K, Irving L, Steinfort DP. Prevalence of incidental pulmonary nodules on computed tomography of the thorax in trauma patients. *Intern Med J*. 2015;45(6):630-633. doi:10.1111/imj.12757.
- Laroche C, Fairbairn I, Moss H, et al. Role of computed tomographic scanning of the thorax prior to bronchoscopy in the investigation of suspected lung cancer. *Thorax*. 2000;55(5):359-363. doi:10.1136/thorax.55.5.359.
- Kim YG, Lee SM, Lee KH, Jang R, Seo JB, Kim N. Optimal matrix size of chest radiographs for computer-aided detection on lung nodule or mass with deep learning. *Eur Radiol*. 2020;30(9):4943-4951. doi:10.1007/s00330-020-06892-9.
- Park S, Lee SM, Lee KH, et al. Deep learning-based detection system for multiclass lesions on chest radiographs: comparison with observer readings. *Eur Radiol*. 2020;30(3):1359-1368. doi:10.1007/s00330-019-06532-x.
- Hernigou J, Cornil F, Poignard A, et al. Thoracic computerised tomography scans in one hundred eighteen orthopaedic patients during the COVID-19 pandemic: identification of chest lesions; added values; help in managing patients; burden on the computerised tomography scan department. *Int Orthop*. 2020;44(8):1571-1580. doi:10.1007/s00264-020-04651-5.
- Erasmus JJ, McAdams HP, Connolly JE. Solitary pulmonary nodules: part II. Evaluation of the indeterminate nodule. *Radiographics*. 2000;20(1):59-66. doi:10.1148/radiographics.20.1.g00ja0259.
- MacMahon H, Austin JH, Gamsu G, et al. Guidelines for management of small pulmonary nodules detected on CT scans: a statement from the Fleischner Society. *Radiology*. 2005;237(2):395-400. doi:10.1148/radiol.2372041887.
- Hoffer FA, Gow K, Flynn PM, Davidoff A. Accuracy of percutaneous lung biopsy for invasive pulmonary aspergillosis. *Pediatr Radiol*. 2001;31(3):144-152. doi:10.1007/s002470000402.
- Nosari A, Anghilieri M, Carrafiello G, et al. Utility of percutaneous lung biopsy for diagnosing filamentous fungal infections in hematologic malignancies. *Haematologica*. 2003;88(12):1405-1409.
- Lange TJ, Kunzendorf F, Pfeifer M, Arzt M, Schulz C. Endobronchial ultrasound-guided transbronchial needle aspiration in routine care: plenty of benign results and follow-up tests. *Int J Clin Pract*. 2012;66(5):438-445. doi:10.1111/j.1742-1241.2012.02907.x.
- Carter BW, Okumura M, Detterbeck FC, Marom EM. Approaching the patient with an anterior mediastinal mass: a guide for radiologists. *J Thorac Oncol*. 2014;9(9 Suppl 2):S110-S118. doi:10.1097/JTO.0000000000000295.
- Laurent F, Montaudon M, Latrabe V, Bégueret H. Percutaneous biopsy in lung cancer. *Eur J Radiol*. 2003;45(1):60-68. doi:10.1016/S0720-048X(02)00286-3.
- Winokur RS, Pua BB, Sullivan BW, Madoff DC. Percutaneous lung biopsy: technique, efficacy, and complications. *Semin Intervent Radiol*. 2013;30(2):121-127. doi:10.1055/s-0033-1342952.
- DiBardino DM, Yarmus LB, Semaan RW. Transthoracic needle biopsy of the lung. *J Thorac Dis*. 2015;7(Suppl 4):S304-S316. doi:10.3978/j.issn.2072-1439.2015.12.16.
- Schneider F, Smith MA, Lane MC, Pantanowitz L, Dacic S, Otori NP. Adequacy of core needle biopsy specimens and fine-needle aspirates for molecular testing of lung adenocarcinomas. *Am J Clin Pathol*. 2015;143(2):193-200. doi:10.1309/AJCPMY8UI7WSFSYY.
- Wang Y, Jiang F, Tan X, Tian P. CT-guided percutaneous transthoracic needle biopsy for paramediastinal and nonparamediastinal lung lesions: diagnostic yield and

- complications in 1484 patients. *Medicine (Baltimore)*. 2016;95(31):e4460. doi:10.1097/MD.0000000000004460.
18. Anzidei M, Sacconi B, Fraioli F, et al. Development of a prediction model and risk score for procedure-related complications in patients undergoing percutaneous computed tomography-guided lung biopsy. *Eur J Cardiothorac Surg*. 2015;48(1):e1-e6. doi:10.1093/ejcts/ezv172.
19. Geraghty PR, Kee ST, McFarlane G, Razavi MK, Sze DY, Dake MD. CT-guided transthoracic needle aspiration biopsy of pulmonary nodules: needle size and pneumothorax rate. *Radiology*. 2003;229(2):475-481. doi:10.1148/radiol.2291020499.
20. Henry M, Arnold T, Harvey J. Pleural Diseases Group, Standards of Care Committee, British Thoracic Society. BTS guidelines for the management of spontaneous pneumothorax. *Thorax*. 2003;58(Suppl 2):ii39-ii52. doi:10.1136/thx.58.suppl\_2.ii39.
21. Hare SS, Gupta A, Goncalves ATC, Souza CA, Matzinger F, Seely JM. Systemic arterial air embolism after percutaneous lung biopsy. *Clin Radiol*. 2011;66(7):589-596. doi:10.1016/j.crad.2011.03.005.



**Copyright:** © 2026 The Author(s). **License:** This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (**CC BY 4.0**), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. For more information, visit, visit <https://creativecommons.org/licenses/by/4.0/>