Neuroscience Insights: Advances in Brain Studies

https://doi.org/10.64978/niabs.2026.11110112

ISSN: 3071-0138

Review Article

Open Access Full Text Article

Eye Colour, Ocular Straylight and Visual Stress: Exploring Potential Associations with Visual Discomfort and Cortical Processing

Michelle Lansdowne[1*]

[1]Chair & Trustee at ISEE (registered Scottish Charitable Incorporated Organisation regulated by the Scottish Charity Regulator (OSCR), UK.

[*]**Correspondence:**

Michelle Lansdowne, Chair & Trustee at ISEE (registered Scottish Charitable Incorporated Organisation regulated by the Scottish Charity Regulator (OSCR), UK.

Received: October 22, 2026 | **Published:** November 11, 2026

How to cite this article:

Lansdowne M. Eye Colour, Ocular Straylight and Visual Stress: Exploring Potential Associations with Visual Discomfort and Cortical Processing. *Neurosci Insights Adv Brain Stud*. 2026;2(4):1-3. doi:10.64978/niabs.2026.11110112.

Abstract

Visual stress is associated with visual discomfort, perceptual distortions and sensitivity to particular visual environments. Although research has investigated the optical characteristics of the eye and the role of visual cortical processing, the potential relationship between eye colour, ocular straylight and the lived experience of visual stress remains largely unexplored. This article proposes that these areas warrant investigation together. Existing research indicates that iris pigmentation may influence ocular straylight, while separate studies suggest that visual discomfort is associated with individual differences in visual processing. However, there is currently insufficient evidence to establish whether these findings are connected to visual stress. This exploratory review examines the rationale for investigating these potential associations, identifies important limitations in current knowledge and proposes a research agenda. It argues that interdisciplinary studies combining objective optical measurements with validated assessments of visual hypersensitivity and self-reported experience could help determine whether eye colour or straylight contributes to particular dimensions of visual discomfort. Rather than proposing a causal explanation, the article seeks to encourage empirical research into a potentially overlooked aspect of visual stress.

Keywords: eye colour; iris pigmentation; ocular straylight; visual stress; visual discomfort; visual hypersensitivity; visual cortex; lived experience.

1. Introduction

Visual stress describes a spectrum of experiences associated with visual discomfort and perceptual disturbance, particularly during reading and exposure to visually demanding environments, which can vary in intensity. Reported symptoms include distortions of text, visual fatigue, headaches and sensitivity to particular patterns and lighting conditions. These experiences vary considerably between individuals and their underlying mechanisms are still being explored.

Research has examined the relationship between visual stress and the processing of particular visual stimuli, including repetitive patterns.[1,2] More recently, researchers have proposed that visual hypersensitivity comprises several dimensions, including sensitivity to brightness, repetitive patterns, strobing and visually intense environments.[7] A broader account of visual discomfort also highlights the potential contribution of differences in cortical processing.[3]

One comparatively under-explored question concerns the possible contribution of individual ocular characteristics. In particular, could eye colour be associated with differences in ocular straylight and might these differences be relevant to visual discomfort experienced by people with visual stress?

The question is not whether eye colour determines visual stress. Rather, it is whether an established relationship between iris pigmentation and aspects of ocular light scatter provides a rationale for investigating possible associations with visual hypersensitivity.

2. Eye Colour and Ocular Straylight

Eye colour primarily reflects differences in iris pigmentation and structure, although pigmentation may have optical implications beyond the visible appearance of the iris.

Van den Berg, Ijspeert and de Waard[10] investigated the relationship between ocular pigmentation and intraocular straylight and their findings indicated that straylight depended on eye colour, particularly at larger scattering angles. They proposed that differences in light transmission through the iris and surrounding ocular wall, together with reflection from the fundus, could contribute to this relationship.

Subsequent research has further examined the relationship between iris colour and visual function. A study by van den Berg et al.[8] Iris colour and visual functions, reported higher intraocular straylight values among participants with light-blue irises than among those with darker irises, suggesting that this difference could be relevant to disability glare in everyday situations.

Straylight occurs when light is scattered within the eye rather than contributing directly to the formation of a clear retinal image. The resulting diffuse luminance can reduce image contrast and impair visual performance, particularly in the presence of bright light sources.[9]

These findings provide a legitimate scientific basis for examining eye colour and straylight together. They do not, however, establish that people with lighter-coloured eyes necessarily experience greater visual discomfort, nor that any difference in straylight is sufficient to produce visual stress. The relationship between objective optical measurements and subjective experience requires dedicated investigation.

3. From Optical Characteristics to Visual Discomfort

Visual stress research suggests that the characteristics of a visual stimulus can influence both perceptual experience and cortical processing. Wilkins[11] discussed how particular visual patterns may provoke discomfort and perceptual distortions in susceptible individuals. More recent work has considered how differences in the processing of visually demanding stimuli might contribute to individual susceptibility.[6]

Evidence concerning discomfort glare is also relevant. Bargary[1] reported differences in cortical activity between individuals with higher and lower sensitivity to discomfort glare. Their findings suggest that the subjective experience of bright light cannot necessarily be explained by optical factors alone.

These research areas raise an important possibility: ocular characteristics and cortical processing may contribute to different stages of the visual experience. Straylight may affect the light distribution reaching the retina, while individual differences in visual processing may influence how the resulting information is experienced.

Whether this interaction occurs in people with visual stress remains unknown. In particular, no conclusion should be drawn that eye colour directly affects cortical excitability or that straylight causes visual stress. These are hypotheses requiring empirical testing.

4. Why This Question Matters

The lived experience of visual stress is heterogeneous: some individuals report difficulties predominantly with text and repetitive patterns, whilst others describe pronounced sensitivity to brightness, glare or visually intense environments. These experiences may overlap without necessarily sharing an identical underlying mechanism.

Price, Sumner and Powell[7] provide a useful framework for investigating this variation by distinguishing several dimensions of visual hypersensitivity. Their findings suggest that future research should avoid treating all visual discomfort as a single phenomenon.

Within this framework, investigating eye colour and straylight could help establish whether optical characteristics are associated with particular forms of visual sensitivity. For example, any association might be more apparent among individuals reporting brightness or glare sensitivity than among those whose principal difficulties involve pattern-related distortions.

This possibility is a research question, not an established finding. Nevertheless, testing it could contribute to a more differentiated understanding of visual stress and help identify whether particular subgroups warrant further investigation.

5. Recommendations for Future Research

An initial observational study could recruit participants with and without self-reported visual stress and assess several variables: objectively classified iris colour, measured ocular straylight, visual acuity, contrast sensitivity and validated measures of visual hypersensitivity.

The analysis should distinguish between different symptom dimensions, including brightness sensitivity, glare discomfort and pattern-related visual stress. Age and relevant ocular characteristics should also be considered, as these may influence straylight independently of iris pigmentation.

A subsequent study could investigate whether straylight measurements are associated with differences in cortical responses to controlled visual stimuli. Such research would require appropriate expertise in vision science and neuroscience, with methods selected and interpreted by suitably qualified researchers.

Importantly, studies should not assume that eye colour is a reliable proxy for straylight or that either characteristic predicts visual stress. Direct measurement is essential. Researchers should also consider alternative explanations and avoid interpreting any observed association as evidence of causation.

The lived experiences of people with visual stress should remain central to this work. Combining objective measurements with participants' descriptions of their symptoms and the environments in which they occur would help establish whether laboratory findings have meaningful relevance to everyday experience.

6. Conclusion

Existing research provides evidence that iris pigmentation can be associated with differences in ocular straylight and that visual discomfort may involve individual differences in cortical processing. However, the relationship between these findings and the lived experience of visual stress has not been established.

Investigating potential associations between eye colour, straylight and visual hypersensitivity could therefore offer a useful avenue for exploratory research. The priority should be to establish whether any measurable relationships exist, whether they differ across dimensions of visual discomfort and whether they have practical significance for affected individuals.

An interdisciplinary approach involving optometry, ophthalmology, vision science, neuroscience and lived-experience research would be particularly valuable. By bringing these perspectives together, researchers could determine whether this proposed connection merits further investigation and contribute to a more comprehensive understanding of visual stress.

References

1. Bargary G, Furlan M, Raynham PJ, Barbur JL, Smith AT. Cortical hyperexcitability and sensitivity to discomfort glare. *Neuropsychologia*. 2015;69:194-200. doi:10.1016/j.neuropsychologia.2015.02.006.
2. Evans BJW, Stevenson SJ. The pattern glare test: a review and determination of normative values. *Ophthalmic Physiol Opt*. 2008;28(4):295-309. doi:10.1111/j.1475-1313.2008.00578.x.
3. Hibbard PB, Allen P, Asher JM et al. A cerebral basis for visual discomfort and visual stress. *Vision*. 2026;10(2):34. doi:10.3390/vision10020034.
4. Huang J, Zhu DC. Visually stressful striped patterns alter human visual cortical functional connectivity. *Hum Brain Mapp*. 2017;38(11):5474-5484. doi:10.1002/hbm.23740.
5. Lansdowne M. Beyond reading: towards a person-centred conceptualisation of visual stress through lived experience and contemporary evidence. *Appl Med Res*. 2026;13(1):283. doi:10.47363/AMR/2026(13)283.
6. Penacchio O, Otazu X, Wilkins AJ, Haigh SM. A mechanistic account of visual discomfort. *Front Neurosci*. 2023;17:1191091. doi.org/10.3389/fnins.2023.1200661.
7. Price A, Sumner P, Powell G. Understanding the subtypes of visual hypersensitivity: four coherent factors and their measurement with the Cardiff Hypersensitivity Scale (CHYPS). *Vision Res*. 2025;233:108610. doi:10.1016/j.visres.2025.108610.
8. van den Berg TJTP, Nischler C, Michael R, et al. Iris color and visual functions. *Graefes Arch Clin Exp Ophthalmol.* 2013;251(1):195-202. doi: 10.1007/s00417-012-2006-8.
9. van den Berg TJTP, Franssen L, Coppens JE. Straylight in the human eye: testing objectivity and optical character of the psychophysical measurement. *Ophthalmic Physiol Opt*. 2009;29(3):345-350. doi:10.1111/j.1475-1313.2009.00638.x.
10. van den Berg TJTP, Ijspeert JK, de Waard PWT. Dependence of intraocular straylight on pigmentation and light transmission through the ocular wall. *Vision Res*. 1991;31(7-8):1361-1367. doi:10.1016/0042-6989(91)90057-C.
11. Wilkins A, Huang J, Cao Y. Visual stress theory and its application to reading and reading tests. *J Res Read*. 2004;27(2):152-162. doi:10.1111/j.1467-9817.2004.00223.x.

Copyright: ©2026 Michelle Lansdowne. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. To view a copy of this licence, visit https://creativecommons.org/licenses/by/4.0/